

OPEN ACCESS

*Correspondence

Ben O. Johnson

Article Received

05/01/2026

Accepted

17/05/2026

Published

06/04/2026

Works Cited

Ben O. Johnson & Auta T. Jonathan, (2026). Computing the Order, Conjugacy Classes and Character Table of the Full Non-Rigid Group of Cyclopropane Chemical Compound Using Wreath Product. *Journal of Current Research and Studies*, 3(2), 25-29.

*COPYRIGHT

© 2026 Ben O. Johnson. This is an open-access article distributed under the terms of the [Creative Commons Attribution License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms

Computing the Order, Conjugacy Classes and Character Table of the Full Non-Rigid Group of Cyclopropane Chemical Compound Using Wreath Product

Ben O. Johnson¹ & Auta T. Jonathan²

^{1,2}Department of Mathematics and Statistics, Federal University, Wukari, Taraba State, Nigeria.

Abstract

The full non-rigid molecules group (f-NRG) theory is seen to be used advantageously to study the internal dynamics of such molecules. In this paper, we determine the order of Cyclopropane and its conjugacy classes. We used computational method to compute the group and its character table and considered the point group of the compound which turned out to be isomorphic to the Wreath Products C_3wrC_2 , where C_n denotes a cyclic group of order n . We found Cyclopropane to have order 18 and 9 conjugacy classes.

Keywords:

Non-Rigid Group, Cyclopropane, WreathProduct, Character Table, Conjugacy Classes

1.0 INTRODUCTION

The application of Group theory concepts to full non-rigid molecules is becoming increasingly relevant and is dense in the research space; see [1] and [2]. Symmetry as a concept is more of mathematics subject and not chemistry. However, the underlying mathematical theory for symmetry and group theory are of great importance in chemistry because they can be used to solve quite plenty of chemistry problems. For example, it can be applied when it comes to the classification of the structures of molecules and crystals, understanding chemical bonding, predicting vibrational spectra and determining the optical activity of compounds.

Cyclopropane exists as a chemical compound with molecular formula as C_3H_6 . Full rigid molecules' symmetry properties are well known and so investigating them is a natural thing to do, though study in this research area is a new in chemistry. [3] presented a number of papers on the determination of character tables of restricted f-NRG of some chemical molecules. For example, Computation of the r-NRG of the triple equivalent methyl rotation in pyramidal trimethylamine with inversion was done by [4]. The authors proved that the r-NRG of this molecule is a group has 648 elements, consists two subgroups of order 324 without inversion. We also have in [5] and [6] where computation of full non-rigid group of some other molecules was carried out. In the same vein, [7] computed full non-rigid group of tetra-ammine platinum (II), trimethylamine, tetraamine platinum (II), etc. which are all examples of C_{2v} and

C_{4v} point groups. Authors such as [8] and [9] proved that molecular structure tetra-amine platinum (II) is isomorphic to the Wreath Product of certain degree. [10] worked on hexamethylbenzene while [11] and [12] studied melamine.

2.0 MATERIALS AND METHODS

Let Ω be a nonempty set, a permutation of Ω is a bijection $\alpha: \Omega \rightarrow \Omega$. We denote the set of all permutations of Ω by S_Ω . When Ω is finite, that is, $\Omega = \{1, 2, \dots, n\}$, we write S_n (the symmetric group of degree n) instead of S_Ω where $|S_n| = n!$ is the number of elements in S_n referred to as the order of the group S_n . Joseph-Louis Lagrange during his time observed permutation as arrangements, that is, as a list i_1, i_2, \dots, i_n with no repetition of any of the elements of Ω . Given an arrangement, i_1, i_2, \dots, i_n , define a function $\alpha: \Omega \rightarrow \Omega$ by $\alpha(j) = i$ for all $j \in \Omega$. Thus, every rearrangement gives a bijection ([13]). The Wreath product of two permutation groups C by D denoted by $W = C \text{ wr } D$ is the semi-direct product of P (a set of mappings) by D , so that, $W = \{(f, d) | f \in P, d \in D\}$, with multiplication in W defined as $(f_1, d_1)(f_2, d_2) = ((f_1, f_2 d_1^{-1}), (d_1, d_2)) \quad \forall f_1, f_2 \in P \text{ and } d_1, d_2 \in D$. W is a special form of permutation group. Understanding the nature of the Wreath products facilitates comprehension of certain types of subgroups of the symmetric groups.

In this paper, we carry out computation of Cyclopropane. We start by first, specifying the algebraic structure of the non-rigid group of these molecules. Then, on the bases of the structure of the group, a useful programming language, namely, "Groups, Algorithms and Programming" (GAP) is applied and the character Tables is computed. The use of this computational group theory package (GAP) is capable of revealing many properties of the groups according to [14] and [15]. We determine the order of the f-NRG of the group and its conjugacy classes. Finally, we compute the character tables of the molecules with the aid of the computational group theory (GAP).

3.0 MAIN RESULT AND DISCUSSION

The Cyclopropane (C_3H_6) ion belongs to the Dihedral groups which are point groups that have an additional C_2 axes that stand perpendicular to the principal axis of the order n . Since there are no other symmetry elements, the point group is of the type D_{3h} .




Figure 1: Cyclopropane Structure.




Figure 2: Cyclopropane Geometry.

Where A = Chemical structure of **Cyclopropane**

B = Geometry of **Cyclopropane**

From figure 1 we have a C_3 principal axis, and three additional C_2 axes, but no other symmetry element. The C_3 axis stands perpendicular to the paper plane, and there are three C_2 axes in the paper plane. Therefore, the symmetry of C_3H_6 has a trigonal bipyramidal shape. The C_3 axis goes through the centre of the atom of the C molecules, and the three C_2 axes go through the three equatorial H atoms. The C_3H_6 symmetry elements, namely C_3 and C_2 axes, are denoted by $C_3 := \langle(1,2,3)\rangle$ and $C_{21} := \langle(4,5)\rangle$, $C_{22} := \langle(6,7)\rangle$, $C_{23} := \langle(8,9)\rangle$ respectively. Therefore, the full symmetry of Cyclopropane as can viewed clearly from figure 2, is: $G = (C_{21} \times C_{22} \times C_{23}) \rtimes C_3$, which we can write in terms of wreath product as $G := \mathbb{Z}_2 \text{ wr } \mathbb{Z}_3$. We used GAP package to get the group as follows:

```
gap>
gap> C3: = Group ((1, 2, 3));
Group ([ (1, 2, 3)])
gap> C2: = Group ((4, 5));
Group ([ (4, 5)])
gap> G: = WreathProduct (C3, C2);
Group ([ (1,2,3), (4,5,6), (1,4) (2,5) (3,6)])
gap> Order (G);
18
gap> CC: = ConjugacyClasses(G);
[ () ^G, (4,5,6) ^G, (4,6,5) ^G, (1,2,3) (4,5,6) ^G, (1,2,3) (4,6,5) ^G, (1,3,2) (4,6,5) ^G,
(1, 4) (2, 5) (3, 6) ^G, (1,4,2,5,3,6) ^G, (1,4,3,6,2,5) ^G]
gap> List (CC, x -> Order (Representative(x)));
[ 1, 3, 3, 3, 3, 3, 2, 6, 6]
gap> Display (CharacterTable(G));
```

Table 1: The Representatives of Conjugacy Classes for Cyclopropane

S/N	Representatives	Size	Name
1.	()	1	1a
2.	(4,5,6)	3	3a
3.	(4,6,5)	3	3b
4.	(1,2,3)(4,5,6)	3	3c
5.	(1,2,3)(4,6,5)	3	3d
6.	(1,3,2)(4,6,5)	3	3e
7.	(1,4)(2,5)(3,6)	2	2a
8.	(1,4,2,5,3,6)	6	6a
9.	(1,4,3,6,2,5)	6	6b

Table 2: Character Table for Cyclopropane

	1a	3a	3b	3c	3d	3e	2a	6a	6b
2P	1a	3b	3a	3e	3d	3c	1a	3c	3e
3P	1a	1a	1a	1a	1a	1a	2a	2a	2a
5P	1a	3b	3a	3e	3d	3c	2a	6b	6a
χ_1	1	1	1	1	1	1	1	1	1
χ_2	1	1	1	1	1	1	-1	-1	-1
χ_3	1	A	\bar{A}	\bar{A}	1	A	-1	-A	$-\bar{A}$
χ_4	1	\bar{A}	A	A	1	\bar{A}	-1	$-\bar{A}$	-A
χ_5	1	A	\bar{A}	\bar{A}	1	A	1	A	\bar{A}
χ_6	1	\bar{A}	-1	A	1	\bar{A}	1	\bar{A}	A
χ_7	2	-1	A	2	-1	2	0	0	0
χ_8	2	\bar{A}	-A	B	-1	\bar{B}	0	0	0
χ_9	2	-A	$-\bar{A}$	\bar{B}	-1	B	0	0	0

Where:

$$A = E(3)^2 = (-1 - \text{Sqrt}(-3))/2 = -1 - b3$$

$$B = 2 * E(3)^2 = -1 - \text{Sqrt}(-3) = -1 + i3$$

Note: Table 1 provides the Representatives of the conjugacy classes of the group while Table 2 gives the Character table. Across the top of the character table is a list of all the symmetry operations in the point group. Down the left side of the character table is a list of irreducible representations. An irreducible representation describes how something (an atom, bond, orbital, etc.) changes when the different symmetry operations are performed. It could stay

the same (1), become inverted (-1), or change completely (0). Representations with A and B labels indicate non-degeneracy. Just below the Table 2 are mathematical functions which can be used to assign orbitals to the irreducible representations.

4.0 CONCLUSIONS

In this work, the symmetry operations of the non-rigid Cyclopropane were studied. The conjugacy classes and the irreducible character tables of the molecules were computed as seen in table 1 and 2 using computational group theory. We found that the group is isomorphic to the Wreath Products $\mathbb{Z}_2 \wr \mathbb{Z}_3$ with order 18 and 9 conjugacy classes where \mathbb{Z}_n are cyclic groups consisting n elements. We used GAP 4.11.1 for our calculations. The character tables obtained give room for the needed classification of wave functions, determination of selection rules, and so on. Thus, symmetry can be applied in the determination of orbital overlap for molecular orbitals more quickly and easily. Note, only orbitals that have the same irreducible representation have the correct symmetry to overlap and form a molecular orbital.

REFERENCES

- 1) Ezra, G. S. (1982). Symmetry Properties of Molecules, Lecture Note in Chemistry 28, Springer, pp. 655-661.
- 2) Maruani, J. and Serre, J. (1983). Symmetries and Properties of Non-Rigid Molecules, (Eds.) Elsevier, Amsterdam, pp.88-92.
- 3) Smeyers, Y. G. (1992). Introduction to group theory for non-rigid molecules. Adv. Quantum Chem., 24: 1-77.
- 4) Smeyers, Y. G. Villa M (2000). A study of the internal dynamics of trimethylamine by means of the non-rigid group theory. J. Math. Chem., 28: 377-388. The GAP Group (2008). GAP Groups, Algorithms, and Programming, Version 4.4.12 (<http://www.gap-system.org/>).
- 5) Balasubramanian, K. (2004a). Nonrigid Group Theory, Tunneling Splittings, and Nuclear Spin Statistics of Water Pentamer: (H₂O)₅. J. Phys. Chem., 108: 5527-5536.
- 6) Balasubramanian, K. (2004b). Group theoretical analysis of vibrational modes and rovibronic levels of extended aromatic C₄₈N₁₂ AZ fullerene. Chem. Phys. Letters, 391: 64-68.
- 7) Ashrafi, A. R. and Hamadani, M. (2003). The full non-rigid group theory for tetraaminoplatinum (II). Croat. Chem. Acta., 76: 299-303.
- 8) Ashrafi, A. R. and Hamadani, M. (2004). Group theory for tetraamine platinum (II) with C_{2v} and C_{4v} point group in the non-rigid system. J. Appl. Math. Comput., 14: 289-303.
- 9) Darafsheh, M. R., Farjami, Y. and Ashrafi, A. R. (2005a). The non-rigid group of tetraamineplatinum (II) as a wreath product. Bull. Chem. Soc. Jpn., 78: 996-1000.
- 10) Darafsheh, M. R., Ashrafi, A. R. and Darafsheh, A. (2005b). Computing the full non-rigid group of tert-butyltetrahydrofuran using wreath product. Int. J. Quantum Chem., 105: 485-492.
- 11) Ashrafi, A. R. and Hamadani, M. (2005). Full non-rigid group theory and symmetry of melamine. J. Iran. Chem. Soc., 2: 135-139.
- 12) Burness, T. C. and Tong-Viet, H. P. (2016). Primitive permutation groups and derangements of prime power order. *Manuscripta Mathematica*. (150)3. Pp. 255–291.
- 13) Karimi, T. E., Moghadam, M. D. G., Farrokhi M. and Aghaei, M. (2011). The full non-rigid group theory for trimethyl borane with C_{3v}, C_{3h} and C_s point groups, J. Argent. Chem. Soc., pp. 67-73.
- 14) Moghani, A., Naghdi, S. and Sorouhesh, M. R. (2010). The Fujita combinatorial enumeration for the non-rigid group of 2, 4- dimethylbenzene. J. Serb. Chem. Soc., 75(1): 91-99.
- 15) Ben O. J., Auta T. J. (2024). Determination of the Conjugacy Classes and Character Table of the Full Non-Rigid Group of Hexachlorocyclopropane Chemical Compound Via Wreath Product of Pair of Permutation Groups. African Journal of Mathematics and Statistics Studies 7(1), 97 103. DOI: 10.52589/AJMST9JSEVAG